

**Table (6):** Relation between outcome & associated fractures among the studied cases.

Variable	Good (n=4)		Excellent (n=8)		$\chi^2$	P
	No	%	No	%		
<b>Associated fractures:</b>						
• No	2	50	8	100		
• Yes:						
-Ipsilateral mid 1/3 clavicle	1	25	0	0	<b>4.80</b>	<b>0.09</b> NS
-Ipsilateral mid 1/3 femur	1	25	0	0		

**Table (7):** Relation between outcome & associated medical conditions among the studied cases.

Variable	Good (n=4)		Excellent (n=8)		$\chi^2$	P
	No	%	No	%		
<b>Associated medical conditions:</b>						
• No	4	100	5	62.5		
• Yes:						
-B.A	0	0	2	25	<b>2</b>	<b>0.37</b> NS
-Type 1 DM	0	0	1	12.5		

**Table (8):** Relation between outcome & time of fracture union and implant removal among the studied cases.

Variable	Good (n=4)	Excellent (n=8)	MW	P
<b>Time: (weeks)</b>				
• Mean $\pm$ SD	6.24 $\pm$ 2.01	5.98 $\pm$ 2.31		
• Median	6	5	<b>0.19</b>	<b>0.86</b> NS
• Range	6 - 10	5 - 10		

**Table (9):** Relation between outcome & difference between the two sides in supination / pronation and among the studied cases.

Variable	Good (n=4)		Excellent (n=8)		$\chi^2$	P
	No	%	No	%		
<b>Had Difference in forearm rotation:</b>						
• No	0	0	5	62.5	<b>4.29</b>	<b>0.03*</b>
• Yes	4	100	3	37.5		

**Table (10):** Relation between outcome & complications among the studied cases.

Variable	Good (n=4)		Excellent (n=8)		$\chi^2$	P
	No	%	No	%		
<b>Complications:</b>						
• No	4	100	5	62.5	<b>2</b>	<b>0.16 NS</b>
• Yes: ( <i>Pin tract infection</i> )	0	0	3	37.5		