

**Table S1.** Relation between laboratory data of studied cases and different subtypes of BAFF gene expression:

<i>BAFF</i>		CC N=8	CT N=22	TT N=10	Test value	P value
<i>At Admission Hb (g/dl)</i>		11.4±0.75 11.5(10.1-12.5)	10.78±1.58 11.1(7.3-14)	10.85±1.71 11.45(7.3-12.7)	1.007 KW	0.604 (NS)
<i>At Admission PLT (x10<sup>3</sup>/mm<sup>3</sup>)</i>		13.75±15.34 9(1-45)	18.23±17.1 13.5(1-60)	17.2±15.7 15(1-42)	0.318 KW	0.853 (NS)
<i>Before discharge Hb (g/dl)</i>		11.54±0.89 11.5(10.1-13)	10.96±1.33 11(8.4-14.2)	10.83±1.41 11.15(7.6-12.7)	1.745 KW	0.418 (NS)
<i>Before discharge PLT(x10<sup>3</sup>/mm<sup>3</sup>)</i>		90.9±133.8 18(4-350)	36.8±49.5 23(2-220)	77.1±82.26 50.5(2-266)	2.070 KW	0.355 (NS)
<i>At 5<sup>th</sup> Month Hb (g/dl)</i>		11.67±0.52 11.7(10.9-12.7)	10.91±1.15 11.05(8.3-12.7)	10.96±1.75 11.25(8-13)	2.824 KW	0.244 (NS)
<i>At 5<sup>th</sup> Month PLT (x10<sup>3</sup>/mm<sup>3</sup>)</i>		120.4±83.2 119(10-236)	63.23±50.82 55(5-205)	116±113 77.5(14-333)	3.270 KW	0.195 (NS)
<i>At 1year Hb (g/dl)</i>		11.13±0.67 10.9(10.4-12)	11.3±1.26 11.35(9.3-13.7)	11.48±1.09 11.55(9-12.9)	1.086 KW	0.581 (NS)
<i>At 1 year PLT (x10<sup>3</sup>/mm<sup>3</sup>)</i>		134.7±119.1 105.5(5-299)	106.5±97.2 80.5(10-391)	146.1±95.28 152(10-300)	1.413 KW	0.493 (NS)
<i>Bone Marrow</i>	<i>Normocellular</i>	5 (62.5%)	8 (36.4%)	8 (80%)	5.650 X2	0.059 (NS)
	<i>Hypercellular</i>	3 (37.5%)	14 (63.6%)	2 (20%)		

KW: Kruscal wallis test, X2: chisquare test, NS: Non significant