

Table (s1): Specimen weight distribution

Mean± SD	798.33±67.53
Median (Range)	800.0 (700-900)

Table (s2): Outcome, result and complication distribution among studied group

		N	%
Hematoma	No	12	100.0
	Yes	0	0.0
Seroma	No	10	83.3
	Yes	2	16.7
Skin necrosis	No	12	100.0
	Yes	0	0.0
Wound dehiscence	No	11	91.7
	Yes	1	8.3
Fat necrosis	No	10	83.3
	Yes	2	16.7
Infection	No	10	83.3
	Yes	2	16.7
Bad scar	No	9	75.0
	Yes	3	25.0
Loss sensation	No	9	75.0
	Yes	3	25.0
Diminished breast milk	No	11	91.7
	Yes	1	8.3
Overall complication	No	9	75.0
	Yes	3	25.0
	Total	12	100.0

Table (s3): Patients satisfaction distribution between groups

		N	%
Patients satisfaction	Not	4	33.3
	Satisfied	8	66.7
	Total	12	100.0

Table (s4): Age, weight and BMI comparison between complicated and not complicated cases

	Complicated (N=3)	Not (N=9)	t	P
Age	39.66±4.72	37.44±2.69	1.039	0.323
Weight	84.33±6.02	86.77±5.91	-0.618	0.551
BMI	28.36±1.0	30.47±2.27	-1.521	0.159

Table (s5): HB pre and post comparison between complicated and not complicated cases

	Complicated (N=3)	Not (N=9)	t	P
HB_PRE	12.53±0.15	12.18±0.52	1.083	0.304
HB_POST	11.93±0.56	11.92±0.65	0.026	0.980

Table (s6): Specimen weight comparison

	Complicated (N=3)	Not (N=9)	t	P
Specimen weight	762.5 ± 91.6	836.6±115.9	1.73	0.078

Table (s7): Association between complication and Clinical and demographic characters

			Overall complication		Total	X ²	P
			Not	Complicated			
Smoking	No	N	8	3	11	0.36	0.54
		%	88.9%	100.0%	91.7%		
	Yes	N	1	0	1		
		%	11.1%	0.0%	8.3%		
DM	No	N	7	2	9	0.14	0.7
		%	77.8%	66.7%	75.0%		
	Yes	N	2	1	3		
		%	22.2%	33.3%	25.0%		
Married	No	N	3	0	3	1.33	0.24
		%	33.3%	0.0%	25.0%		
	Yes	N	6	3	9		
		%	66.7%	100.0%	75.0%		
Have children	No	N	5	0	5	2.85	0.091
		%	55.6%	0.0%	41.7%		
	Yes	N	4	3	7		
		%	44.4%	100.0%	58.3%		
Breast feeding	No	N	5	0	5	2.85	0.091
		%	55.6%	0.0%	41.7%		
	Yes	N	4	3	7		
		%	44.4%	100.0%	58.3%		
Total		N	9	3	12		
		%	100.0%	100.0%	100.0%		